



TITLE:

# Narrow-band imaging for the head and neck region and the upper gastrointestinal tract.

AUTHOR(S):

Kikuchi, Osamu; Ezoe, Yasumasa; Morita, Shuko; Horimatsu, Takahiro; Muto, Manabu

---

CITATION:

Kikuchi, Osamu ...[et al]. Narrow-band imaging for the head and neck region and the upper gastrointestinal tract.. Japanese journal of clinical oncology 2013, 43(5): 458-465

ISSUE DATE:

2013-05

URL:

<http://hdl.handle.net/2433/189840>

RIGHT:

This is a pre-copyedited, author-produced PDF of an article accepted for publication in "Japanese journal of clinical oncology" following peer review. The version of record "Osamu Kikuchi, Yasumasa Ezoe, Shuko Morita, Takahiro Horimatsu, and Manabu Muto; Narrow-band Imaging for the Head and Neck Region and the Upper Gastrointestinal Tract; Jpn. J. Clin. Oncol. (2013) 43 (5): 458-465 doi:10.1093/jjco/ht042" is available online at: <http://jjco.oxfordjournals.org/content/43/5/458>; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

# **Narrow band imaging for head and neck region and upper gastrointestinal tract**

**Osamu Kikuchi<sup>1</sup>, Yasumasa Ezoe<sup>2</sup>, Shuko Morita<sup>1</sup>, Takahiro Horimatsu<sup>1</sup>, Manabu Muto<sup>3</sup>**

**1, Department of Gastroenterology and Hepatology,**

**2, Department of Multidisciplinary Cancer Treatment**

**3, Department of Therapeutic Oncology**

**Kyoto University Graduate School of Medicine, 54 Shogoin kawaharacho,**

**Sakyoku, Kyoto, 606-8507, Japan**

**Correspondence to: Manabu Muto**

**Department of Therapeutic Oncology, Kyoto University Hospital**

**54 Shogoin kawaharacho, Sakyoku, Kyoto, 606-8507, Japan**

**Tel: +81-75-751-4319, Fax: +81-75-751-4303**

**E-mail: [mmuto@kuhp.kyoto-u.ac.jp](mailto:mmuto@kuhp.kyoto-u.ac.jp)**

## **Abstract**

Endoscopy is essential for diagnosis and treatment of cancers derived from gastrointestinal tract. However, conventional white light image has technical limitation in detecting small or superficial lesions. Narrow band imaging, especially with magnification, allows visualize a microstructure pattern and microvascular patterns in the mucosal surface. These technical breakthroughs enable endoscopists to easily detect small pre-neoplastic and neoplastic lesions and to make differential diagnosis of these lesions. Appropriate diagnosis with NBI contributes to minimally invasive endoscopic resection.

## **Mini-abstract**

Narrow-band imaging is discussed, which allows endoscopists to easily diagnose small lesions and to make differential diagnosis of these lesions. Appropriate diagnosis with NBI contributes to minimally invasive treatments.

## ***Key words:***

endoscopy, narrow band imaging, early detection, differential diagnosis

## NARROW BAND IMAGING

Narrow band imaging (NBI) is an innovative optical technology that allows the distinct visualization of microsurface patterns and microvascular patterns on the mucosal surface (1-3). The NBI system uses narrow-band illumination created with optical interference filters that generate 415 nm and 540 nm wavelengths, corresponding to the peaks of absorption of hemoglobin. Therefore, thin blood vessels, such as capillaries, in the epithelium or mucosal layer can be seen more distinctly than in a conventional white-light image (WLI) (Figure 1).

Currently, two types of image reconstruction systems are used for endoscopic imaging: a red–green–blue (RGB) time sequential illumination system with a monochrome charge-coupled device (CCD) and white-light illumination with a color chip CCD. The NBI system is applicable to both systems by placing the narrow-band light filter in front of the light source. NBI can provide the same clinical benefits with both illumination systems (Table 1), although the color reproduction and the image resolution are somewhat different in the two systems (4).



## HEAD AND NECK REGION

### HEAD AND NECK CANCER

Lugol chromoendoscopy is the standard method for detecting early squamous cell carcinoma (SCC) of the esophagus. However, Lugol dye solution cannot be applied to the oropharynx or hypopharynx because of the risk of aspiration. Moreover, the image resolution of rhinolaryngoscopy does not effectively identify superficial neoplastic lesions in the head and neck region. Therefore, the early detection of cancers in the oropharynx and hypopharynx has been difficult. This is partly attributable to the technological limitations in mounting a high-resolution CCD on the tip of a rhinolaryngoscope.

Muto *et al.* first reported the utility of NBI combined with magnifying endoscopy (Q240Z, Olympus Medical Systems, Tokyo, Japan) in the identification of superficial SCCs in the head and neck region (5). Compared with WLI, NBI significantly improved the visualization of the cancerous lesions by enhancing the contrast between the cancerous lesion and the background nonneoplastic epithelium, and by the clear magnification of the microvascular architecture (6). Muto *et al.* reported that the well-demarcated brownish areas observed under NBI and the microvascular

irregularities visible under magnification with NBI were useful indicators of cancerous lesions in the head and neck region (1). In the multicenter prospective randomized study, NBI is revealed to be superior to WLI in the detection and differential diagnosis for superficial head and neck cancer (7).

Watanabe et al. reported that the NBI rhinolaryngoscope (ENF-V2, Olympus Medical Systems) with a color-chip light source (CLV-160B, Olympus Medical Systems) improved the diagnostic accuracy, and the negative predictive values for superficial lesions in the oropharynx and hypopharynx compared with those of conventional WLI (8, 9). However, there is still a critical difference in the image qualities of CCDs between the gastrointestinal endoscopy and that the rhinolaryngoscope.

Ugumori et al. prospectively compared the images taken with a color-chip-based rhinolaryngoscope and those taken with an RGB-sequential-system-based high-resolution gastrointestinal endoscope (10). Whereas the conventional white-light rhinolaryngoscope identified well-demarcated line between the neoplastic and nonneoplastic lesions in only 10% (5/51) of cases and microvascular irregularities in only 27% (14/51), the NBI rhinolaryngoscope identified these in 63% (32/51) and 94% (49/51) of cases, respectively. These results indicate that even with a rhinolaryngoscope, NBI can improve the visualization of epithelial neoplasms of the head and neck region.

When combined with a high-definition television camera (HDTV), the effectiveness of NBI is improved in terms of both its sensitivity and specificity.(11)

NBI is also reportedly useful in detecting metachronous SCC after treatment for esophageal SCC (chemoradiotherapy, radiation therapy, or surgery), and unknown primary SCC of the neck, and adenoid hypertrophy (12-18) (Table 1).

The early detection of cancer in this region increases the possibility of minimally invasive surgery, including endoscopic mucosal resection and endoscopic submucosal dissection methods (19, 20). The potential advantages to patients resulting from an early diagnosis, with the preservation of organ and tissue functions, are obvious.

## **Esophagus**

### **ESOPHAGEAL SQUAMOUS CELL CARCINOMA**

Although the early detection of cancer offers the best prognosis, many esophageal SCCs (ESCCs) are still detected at a late stage, with a consequently poor prognosis. One reason is that the early detection of ESCC is difficult using conventional WLI endoscopy because it cannot identify the morphological changes of superficial ESCC. Although Lugol chromoendoscopy is the sensitive method for the detection of early

superficial ESCC (Figure1 A, B), iodine is an irritant and causes unpleasant reactions, such as pain, discomfort and sometimes allergic reaction. In contrast, NBI is less invasive than Lugol chromoendoscopy and enhances the clarity of the intrapapillary capillary loop (IPCL) patterns beneath the epithelium (5, 21, 22), so it is expected to replace Lugol chromoendoscopy in this role (Figure1 C,D).

Using an ultrathin endoscope (5 mm in diameter at the distal end; XP260N, Olympus Medical Systems), Lee et al. reported the utility of NBI in the detection and accurate diagnosis of ESCC (23). The sensitivity of NBI was significantly better than that of conventional WLI. The specificity and positive predictive value of NBI were also better than those of Lugol chromoendoscopy, whereas their diagnostic accuracy and negative predictive value were similar. These results suggest that, even when an ultrathin endoscope is used, NBI is the best tool for screening for superficial esophageal neoplasms, as in the head and neck region.

In a multicenter prospective randomized study (7), NBI with a standard-diameter endoscope showed approximately twofold greater sensitivity than WLI. Furthermore, most of the Lugol-voiding lesions overlooked by NBI endoscopy were low grade intraepithelial neoplasia or lesions without atypical findings (24).

In 2011, a new classification of magnified endoscopy for superficial ESCC was

proposed by Japan Esophageal Society (2d5), which allows the differential diagnosis of ESCC, intraepithelial neoplasia, and inflammation. This classification is expected to simplify the diagnosis and evaluation of the depth of invasion of superficial ESCCs.

### **GASTROESOPHAGEAL REFLUX DISEASE (GERD)**

GERD is defined by the presence of reflux esophagitis. When it causes reflux symptoms (chest pain, heartburn, discomfort, etc.), the patient's quality of life is adversely affected (26). Moreover, a significant number of patients with GERD symptoms show no endoscopic signs of esophagitis. This condition is described as “nonerosive reflux disease” (NERD). Many NERD patients show minimal endoscopic findings, such as a whitish or reddish edematous change or erosion that is not regarded as a mucosal break (27). These minimal changes are potentially related to various GERD symptoms (28). However, the interobserver agreement when NERD is diagnosed with conventional WLI is reportedly too low to support the clinical significance of this technique (29). In contrast, NBI is expected to overcome this limitation, because it allows the visualization of the superficial and slight findings attributable to NERD, which cannot be seen with conventional WLI.

Lee et al. reported that the intraobserver and interobserver consistencies in grading

esophagitis improved when NBI was used instead of WLI (30). Sharma et al. reported a feasibility study of magnified endoscopy with NBI in patients with GERD (31). They showed that increased numbers and the dilatation of IPCLs were the best predictors of a diagnosis of GERD, with moderate to high interobserver agreement.

### **BARRETT'S ESOPHAGUS AND CANCER**

The incidence of esophageal adenocarcinoma is increasing in Western countries (32) and Barrett's esophagus (BE) is a precursor lesion of this malignancy. Surveillance of BE using WLI with random four-quadrant biopsies is the accepted practice and is recommended by the American Gastroenterological Association statement (33). Sharma et al. (34) showed in a randomized, controlled, international, crossover trial that the success of NBI in detecting intestinal metaplasia did not differ from that of the currently accepted practice of random biopsies, but required significantly fewer biopsies.

Because esophageal adenocarcinoma has a poor prognosis when detected at an advanced stage, endoscopic surveillance is recommended to detect high-grade dysplasia and mucosal neoplasia in patients with BE. However, it is difficult to identify these lesions with conventional WLI. NBI with magnifying endoscopy allows us to visualize the details of the mucosal microsurface pattern and the microvascular pattern without

additional equipment or dye solutions (35).

Hamamoto et al. first reported that NBI could better visualize the esophagogastric junction, net-like capillary vessels, and columnar-lined esophagus (seen in BE) than conventional WLI (36). Kara et al. reported that indigo carmine chromoendoscopy and NBI were similarly effective in the diagnosis of high-grade dysplasia or early cancer in BE (37). Wolfsen et al. (38) reported that high-resolution NBI can detect dysplastic lesions more efficiently, with fewer biopsy samples, than standard-resolution WLI. Singh et al. (39) reported that NBI with magnification is superior to WLI with magnification in the prediction of histology in BE.

A recent meta-analysis (40) that included 446 patients with 2194 lesions reported that NBI with magnification shows high diagnostic precision in detecting high-grade dysplasia, with a sensitivity and specificity of 96% and 94%, respectively.

## **Stomach**

### **DETECTION OF GASTRIC NEOPLASM BY NBI**

In the stomach, NBI has been considered to be used with magnification for detailed examinations. Because the light intensity under the NBI filter is low, a non-magnified image becomes dark compared with that produced under WLI. Furthermore, because

the image becomes noisy with the electrical enhancement used to keep the endoscopic image bright, it is insufficient to observe the wide area of the stomach. There is also, as yet, no evidence that NBI is superior to WLI in detecting early gastric neoplasms. To overcome these limitations, much brighter NBI system with higher resolution will be commercially available when this review is published. Then, the evidence of other clinical benefit of NBI such as detection will be expected in future.

## **DIFFERENTIAL DIAGNOSIS OF GASTRIC CANCER**

Yao et al. originally reported unique magnifying endoscopic findings of gastric cancer in 2002 (41). This marked the beginning of the era of using magnifying endoscopy for the diagnosis of gastric cancer. The utility of magnifying endoscopic observations combined with WLI for the differential diagnosis of flat or slightly depressed gastric cancers and nonneoplastic lesions, such as gastritis, has been reported. NBI combined with magnifying endoscopy (magnifying NBI) provides better visualization of the mucosal surface and microvascular architecture than magnifying WLI (42). Several reports have compared the diagnostic yield of magnifying NBI with that of magnifying or nonmagnifying WLI in distinguishing small gastric cancers from the flat or depressed benign lesions caused by chronic gastritis (43-45). However, all those reports had some



limitations: they were performed at only one institution, evaluated stored images and did not involve real-time assessment, or included gastric lesions with a definite pathological diagnosis. To overcome these limitations, Ezoe et al. performed a multicenter, prospective, randomized, controlled trial that targeted newly detected, undiagnosed lesions to compare and evaluate the diagnostic yields of magnifying NBI and conventional WLI. The trial revealed that magnifying NBI, especially after nonmagnifying WLI, showed an extremely high diagnostic performance (46).

These lines of evidence suggest that magnifying NBI is currently one of the standard endoscopic modalities in the differential diagnosis of gastric cancers.

## **DETERMINATION OF THE LATERAL EXTENT OF GASTRIC CANCER**

To achieve the complete resection of a mucosal gastric cancer with endoscopic resection, an accurate diagnosis of the extent of the tumor is required. By clearly visualizing the microvascular architecture and the microsurface structure inside and outside the lesion, magnifying NBI can distinguish the cancer margins from the surrounding benign mucosa, so it is expected to be useful for delineating the extent of a gastric tumor. In 2004, Sumiyama et al. retrospectively described the feasibility of NBI for the guidance of *en bloc* endoscopic resection when combined with a multibending endoscope, but did

not perform a formal evaluation (47). Kadowaki et al. compared the utility of magnifying NBI and magnifying WLI in recognizing gastric cancer demarcation. They also reported that magnifying NBI is more useful when it is combined with acetic acid (48). Kiyotoki et al. (49) and Nagahama et al. (50) reported the superiority of magnifying NBI to chromoendoscopy for determining the lateral extent of early gastric cancer. These lines of evidence suggest that magnifying NBI can be a useful modality for determining the lateral extent of gastric cancer. However, it must be emphasized that it is still difficult to accurately define the tumor margin in undifferentiated gastric cancers; the successful delineation rate was 0% for undifferentiated cancers in one study (50). Because undifferentiated gastric cancers often spread subepithelially and are covered with nonneoplastic foveolar epithelium, observation of the mucosal surface by NBI is not useful for determining the tumor margin of this type of gastric cancer. Therefore, it is necessary to take biopsy specimens of the surrounding mucosa to define the extent of an otherwise undetectable tumor in undifferentiated gastric cancers.

## **PREDICTION OF THE HISTOLOGICAL TYPE OF GASTRIC CANCER**

Nakayoshi et al. reported that the different microvascular patterns detected with magnifying NBI images are useful in predicting the histological type of a superficial

gastric cancer (51). Differentiated adenocarcinomas display a “fine network pattern,” and undifferentiated adenocarcinomas display a “corkscrew pattern” in their microvascular structures (Figure 2). Yoshida et al. reported that a “nonstructural pattern” appeared to be a useful marker of undifferentiated superficial gastric cancers (52).

Although these studies have indicated the utility of magnifying NBI in the prediction of the histopathological type of a gastric cancer, its reliability must be validated in a large-scale prospective study. Moreover, even if magnifying NBI can predict the histological type of a cancerous lesion, histological confirmation by biopsy is required at this time. However, the prediction of histological type could be useful to the endoscopists when selecting the site of a biopsy in a lesion because gastric cancers are usually heterogeneous.

## **DIAGNOSIS OF THE TUMOR DEPTH OF GASTRIC CANCER**

In contrast to ESCC, there is no evidence that NBI, with or without magnifying endoscopy, can predict the depth of tumor invasion in a patient with gastric cancer.

## **DIAGNOSIS OF GASTRIC ADENOMA**

Because most gastric adenomas form protruded lesions, the differential diagnosis of

protruded gastric cancer and protruded adenoma is sometimes difficult (53, 54). Yao et al. reported that the characteristic finding of magnifying NBI, a white opaque substance (Figure 3), is a relevant sign for differentiating protruded adenomas from protruded cancers (55). Tsuji et al. also reported that the presence of an irregular microvascular pattern or irregular microsurface pattern with a demarcation line between the lesion and the surrounding area under magnifying NBI is useful in distinguishing cancers from adenomas (56). Maki et al. reported that magnifying NBI appears to be useful in differentiating between cancerous and adenomatous superficial elevated lesions of the stomach with significantly higher sensitivity and accuracy (57). In contrast, depressed-type adenoma is rare, although it is clinically important because it has greater malignant potential than protruded adenoma (58). Tamai et al. reported that depressed-type adenomas display a regular ultrafine pattern, in which the network of microvessels is composed of small and regular circles, which differs from the irregular fine network pattern of well-differentiated gastric cancers (59).

These reports indicate that magnifying NBI should be a useful modality for the accurate diagnosis of gastric adenoma.

## Duodenum

Small duodenal ampullary tumors are treated by surgical resection or endoscopic resection. However, the lateral margin must be precisely assessed before curative endoscopic resection. Uchiyama et al. reported that magnifying NBI with a direct frontal-view magnifying endoscope can predict the histological characteristics of ampullary lesions by detecting abnormal vessels and microsurface patterns (60). Itoi et al. reported that NBI with a conventional duodenoscope, with no magnifying capacity, allowed better visualization of the tumor margin than indigo carmine chromoendoscopy (61). However, these studies included only a small number of cases, so further studies with a sufficient number of patients are required to evaluate the usefulness of NBI for duodenal tumors.

## Conclusion

NBI is now useful endoscopic modality for head and neck region and the upper gastrointestinal tract. It helps the endoscopists to do early detection and accurate diagnosis for the head and neck neoplasia and disease in the upper gastrointestinal diseases. Furthermore, it provides the many chance to do minimally invasive treatment and improves the patients' survival and quality of life. Then, standard education

program of NBI in clinical practice will be needed.

## References

1. Muto M, Katada C, Sano Y, Yoshida S. Narrow band imaging: a new diagnostic approach to visualize angiogenesis in superficial neoplasia. Clin Gastroenterol Hepatol. 2005;3:S16-20.
2. Gono K, Obi T, Yamaguchi M, et al. Appearance of enhanced tissue features in narrow-band endoscopic imaging. J Biomed Opt. 2004;9:568-77.
3. Gono K, Yamazaki K, Doguchi N, et al. Endoscopic Observation of Tissue by Narrowband Illumination. Optical Review. 2003;10:211-5.
4. Rey JF, Tanaka S, Lambert R, Tajiri H. Evaluation of the clinical outcomes associated with EXERA II and LUCERA endoscopes. Dig Endosc. 2009;21(Suppl 1):S113-20.
5. Muto M, Nakane M, Katada C, et al. Squamous cell carcinoma in situ at oropharyngeal and hypopharyngeal mucosal sites. Cancer. 2004;101:1375-81.
6. Muto M, Ugumori T, Sano YI, Ohtsu A, Yoshida S. Narrow-band imaging combined with magnified endoscopy for cancer at the head and neck region. Dig Endosc. 2005;17:S23-S4.

7. Muto M, Minashi K, Yano T, et al. Early detection of superficial squamous cell carcinoma in the head and neck region and esophagus by narrow band imaging: a multicenter randomized controlled trial. *J Clin Oncol.* 2010;28:1566-72.
8. Watanabe A, Tsujie H, Taniguchi M, Hosokawa M, Fujita M, Sasaki S. Laryngoscopic detection of pharyngeal carcinoma in situ with narrowband imaging. *Laryngoscope.* 2006;116:650-4.
9. Watanabe A, Taniguchi M, Tsujie H, Hosokawa M, Fujita M, Sasaki S. The value of narrow band imaging endoscope for early head and neck cancers. *Otolaryngol Head Neck Surg.* 2008;138:446-51.
10. Ugumori T, Muto M, Hayashi R, Hayashi T, Kishimoto S. Prospective study of early detection of pharyngeal superficial carcinoma with the narrowband imaging laryngoscope. *Head Neck.* 2009;31:189-94.
11. Piazza C, Cocco D, De Benedetto L, Del Bon F, Nicolai P, Peretti G. Narrow band imaging and high definition television in the assessment of laryngeal cancer: a prospective study on 279 patients. *Eur Arch Otorhinolaryngol.* 2010;267:409-14.
12. Lin YC, Watanabe A, Chen WC, Lee KF, Lee IL, Wang WH.



Narrowband imaging for early detection of malignant tumors and radiation effect after treatment of head and neck cancer. *Arch Otolaryngol Head Neck Surg.* 2010;136:234-9.

13. Piazza C, Cocco D, De Benedetto L, Del Bon F, Nicolai P, Peretti G. Role of narrow-band imaging and high-definition television in the surveillance of head and neck squamous cell cancer after chemo- and/or radiotherapy. *Eur Arch Otorhinolaryngol.* 2010;267:1423-8.

14. Nonaka S, Saito Y, Oda I, Kozu T, Saito D. Narrow-band imaging endoscopy with magnification is useful for detecting metachronous superficial pharyngeal cancer in patients with esophageal squamous cell carcinoma. *J Gastroenterol Hepatol.* 2010;25:264-9.

15. Piazza C, Cocco D, Del Bon F, et al. Narrow band imaging and high definition television in evaluation of oral and oropharyngeal squamous cell cancer: a prospective study. *Oral Oncology.* 2010;46:307-10.

16. Wang WH, Lin YC, Weng HH, Lee KF. Narrow-band imaging for diagnosing adenoid hypertrophy in adults: a simplified grading and histologic correlation. *Laryngoscope.* 2011;121:965-70.

17. Chu PY, Tsai TL, Tai SK, Chang SY. Effectiveness of narrow band

imaging in patients with oral squamous cell carcinoma after treatment.

Head Neck. 2012;34:155-61.

18. Masaki T, Katada C, Nakayama M, et al. Usefulness and pitfall of Narrow band imaging combined with magnifying endoscopy for detecting an unknown head and neck primary site with cervical lymph node metastasis.

Auris, nasus, larynx. 2012;39:502-6.

19. Muto M, Satake H, Yano T, et al. Long-term outcome of transoral organ-preserving pharyngeal endoscopic resection for superficial pharyngeal cancer. Gastrointest Endosc. 2011;74:477-84.

20. Shimizu Y, Yamamoto J, Kato M, et al. Endoscopic submucosal dissection for treatment of early stage hypopharyngeal carcinoma.

Gastrointest Endosc. 2006;64:255-9.

21. Arima M, Tada M, Arima H. Evaluation of microvascular patterns of superficial esophageal cancers by magnifying endoscopy. Esophagus.

2005;2:191-7.

22. Yoshida T, Inoue H, Usui S, Satodate H, Fukami N, Kudo SE.

Narrow-band imaging system with magnifying endoscopy for superficial esophageal lesions. Gastrointest Endosc. 2004;59:288-95.

23. Lee YC, Wang CP, Chen CC, et al. Transnasal endoscopy with narrow-band imaging and Lugol staining to screen patients with head and neck cancer whose condition limits oral intubation with standard endoscope (with video). *Gastrointest Endosc.* 2009;69:408-17.
24. Takenaka R, Kawahara Y, Okada H, et al. Narrow-band imaging provides reliable screening for esophageal malignancy in patients with head and neck cancers. *Am J Gastroenterol.* 2009;104:2942-8.
25. Oyama T, Monma K. Workshop 1: A new classification of magnified endoscopy for superficial esophageal squamous cell carcinoma, in *Summaries from the 65th Annual Meeting of the Japan Esophageal Society on September 26, 2011, Sendai.* *Esophagus.* 2011;8:251.
26. Dent J, Armstrong D, Delaney B, Moayyedi P, Talley NJ, Vakil N. Symptom evaluation in reflux disease: workshop background, processes, terminology, recommendations, and discussion outputs. *Gut.* 2004;53(Suppl 4):iv1-24.
27. Nakamura T, Shirakawa K, Masuyama H, Sugaya H, Hiraishi H, Terano A. Minimal change oesophagitis: a disease with characteristic differences to erosive oesophagitis. *Aliment Pharmacol Ther.* 2005;21(Suppl

2):19-26.

28. Lee JH, Kim N, Chung IK, et al. Clinical significance of minimal change lesions of the esophagus in a healthy Korean population: a nationwide multi-center prospective study. *J Gastroenterol Hepatol*. 2008;23:1153-7.

29. Amano Y, Ishimura N, Furuta K, et al. Interobserver agreement on classifying endoscopic diagnoses of nonerosive esophagitis. *Endoscopy*. 2006;38:1032-5.

30. Lee YC, Lin JT, Chiu HM, et al. Intraobserver and interobserver consistency for grading esophagitis with narrow-band imaging. *Gastrointest Endosc*. 2007;66:230-6.

31. Sharma P, Wani S, Bansal A, et al. A feasibility trial of narrow band imaging endoscopy in patients with gastroesophageal reflux disease. *Gastroenterology*. 2007;133:454-64.

32. Pera M, Manterola C, Vidal O, Grande L. Epidemiology of esophageal adenocarcinoma. *J Surg Oncol*. 2005;92:151-9.

33. Spechler SJ, Sharma P, Souza RF, Inadomi JM, Shaheen NJ. American Gastroenterological Association medical position statement on the management of Barrett's esophagus. *Gastroenterology*. 2011;140:1084-91.

34. Sharma P, Hawes RH, Bansal A, et al. Standard endoscopy with random biopsies versus narrow band imaging targeted biopsies in Barrett's oesophagus: a prospective, international, randomised controlled trial. *Gut*. 2012;published online: 7 February 2012. doi:10.1136/gutjnl-2011-300962.
35. Goda K, Tajiri H, Ikegami M, Urashima M, Nakayoshi T, Kaise M. Usefulness of magnifying endoscopy with narrow band imaging for the detection of specialized intestinal metaplasia in columnar-lined esophagus and Barrett's adenocarcinoma. *Gastrointest Endosc*. 2007;65:36-46.
36. Hamamoto Y, Endo T, Nosho K, Arimura Y, Sato M, Imai K. Usefulness of narrow-band imaging endoscopy for diagnosis of Barrett's esophagus. *J Gastroenterol*. 2004;39:14-20.
37. Kara MA, Peters FP, Rosmolen WD, et al. High-resolution endoscopy plus chromoendoscopy or narrow-band imaging in Barrett's esophagus: a prospective randomized crossover study. *Endoscopy*. 2005;37:929-36.
38. Wolfsen HC, Crook JE, Krishna M, et al. Prospective, controlled tandem endoscopy study of narrow band imaging for dysplasia detection in Barrett's Esophagus. *Gastroenterology*. 2008;135:24-31.
39. Singh R, Karageorgiou H, Owen V, et al. Comparison of high-resolution

magnification narrow-band imaging and white-light endoscopy in the prediction of histology in Barrett's oesophagus. *Scandinavian journal of gastroenterology*. 2009;44:85-92.

40. Mannath J, Subramanian V, Hawkey CJ, Ragunath K. Narrow band imaging for characterization of high grade dysplasia and specialized intestinal metaplasia in Barrett's esophagus: a meta-analysis. *Endoscopy*. 2010;42:351-9.

41. Yao K, Oishi T, Matsui T, Yao T, Iwashita A. Novel magnified endoscopic findings of microvascular architecture in intramucosal gastric cancer. *Gastrointest Endosc*. 2002;56:279-84.

42. Yao K, Iwashita A, Tanabe H, et al. Novel zoom endoscopy technique for diagnosis of small flat gastric cancer: a prospective, blind study. *Clin Gastroenterol Hepatol*. 2007;5:869-78.

43. Kaise M, Kato M, Urashima M, et al. Magnifying endoscopy combined with narrow-band imaging for differential diagnosis of superficial depressed gastric lesions. *Endoscopy*. 2009;41:310-5.

44. Ezoe Y, Muto M, Horimatsu T, et al. Magnifying narrow-band imaging versus magnifying white-light imaging for the differential diagnosis of

gastric small depressive lesions: a prospective study. *Gastrointest Endosc.*

2010;71:477-84.

45. Kato M, Kaise M, Yonezawa J, et al. Magnifying endoscopy with narrow-band imaging achieves superior accuracy in the differential diagnosis of superficial gastric lesions identified with white-light endoscopy: a prospective study. *Gastrointest Endosc.* 2010;72:523-9.

46. Ezoe Y, Muto M, Uedo N, et al. Magnifying narrowband imaging is more accurate than conventional white-light imaging in diagnosis of gastric mucosal cancer. *Gastroenterology.* 2011;141:2017-25.

47. Sumiyama K, Kaise M, Nakayoshi T, et al. Combined use of a magnifying endoscope with a narrow band imaging system and a multibending endoscope for en bloc EMR of early stage gastric cancer. *Gastrointest Endosc.* 2004;60:79-84.

48. Kadowaki S, Tanaka K, Toyoda H, et al. Ease of early gastric cancer demarcation recognition: a comparison of four magnifying endoscopy methods. *J Gastroenterol Hepatol.* 2009;24:1625-30.

49. Kiyotoki S, Nishikawa J, Satake M, et al. Usefulness of magnifying endoscopy with narrow-band imaging for determining gastric tumor margin.

J Gastroenterol Hepatol. 2010;25:1636-41.

50. Nagahama T, Yao K, Maki S, et al. Usefulness of magnifying endoscopy with narrow-band imaging for determining the horizontal extent of early gastric cancer when there is an unclear margin by chromoendoscopy (with video). Gastrointest Endosc. 2011;74:1259-67.

51. Nakayoshi T, Tajiri H, Matsuda K, Kaise M, Ikegami M, Sasaki H. Magnifying endoscopy combined with narrow band imaging system for early gastric cancer: correlation of vascular pattern with histopathology (including video). Endoscopy. 2004;36:1080-4.

52. Yoshida T, Kawachi H, Sasajima K, Shiokawa A, Kudo SE. The clinical meaning of a nonstructural pattern in early gastric cancer on magnifying endoscopy. Gastrointest Endosc. 2005;62:48-54.

53. Kamiya T, Morishita T, Asakura H, Miura S, Munakata Y, Tsuchiya M. Long-term follow-up study on gastric adenoma and its relation to gastric protruded carcinoma. Cancer. 1982;50:2496-503.

54. Kato M, Nishida T, Tsutsui S, et al. Endoscopic submucosal dissection as a treatment for gastric noninvasive neoplasia: a multicenter study by Osaka University ESD Study Group. J Gastroenterol. 2011;46:325-31.



55. Yao K, Iwashita A, Tanabe H, et al. White opaque substance within superficial elevated gastric neoplasia as visualized by magnification endoscopy with narrow-band imaging: a new optical sign for differentiating between adenoma and carcinoma. *Gastrointest Endosc.* 2008;68:574-80.
56. Tsuji Y, Ohata K, Sekiguchi M, et al. Magnifying endoscopy with narrow-band imaging helps determine the management of gastric adenomas. *Gastric cancer.* 2012;published online: 18 January 2012. doi: 10.1007/s10120-011-0133-2.
57. Maki S, Yao K, Nagahama T, et al. Magnifying endoscopy with narrow-band imaging is useful in the differential diagnosis between low-grade adenoma and early cancer of superficial elevated gastric lesions. *Gastric Cancer.* 2012;published online: 18 May 2012. doi:10.1007/s10120-012-0160-7.
58. Nakamura K, Sakaguchi H, Enjoji M. Depressed adenoma of the stomach. *Cancer.* 1988;62:2197-202.
59. Tamai N, Kaise M, Nakayoshi T, et al. Clinical and endoscopic characterization of depressed gastric adenoma. *Endoscopy.* 2006;38:391-4.
60. Uchiyama Y, Imazu H, Kakutani H, et al. New approach to diagnosing

ampullary tumors by magnifying endoscopy combined with a narrow-band imaging system. *J Gastroenterol.* 2006;41:483-90.

61. Itoi T, Tsuji S, Sofuni A, et al. A novel approach emphasizing preoperative margin enhancement of tumor of the major duodenal papilla with narrow-band imaging in comparison to indigo carmine chromoendoscopy (with videos). *Gastrointest Endosc.* 2009;69:136-41.

Table1 Advantage of NBI in contrast to WLI or CE in the clinical practice in the head and neck region and the upper gastrointestinal tract.

Site	Category	Study aim	Evidence	Magnification	Study design	Diagnosis	Reference
Head and neck	Squamous cell carcinoma	Detection	NBI > WLI (accuracy: 86.7 vs. 62.9%)	+	Prospective RCT	On site	7
		Differential diagnosis	NBI > WLI	+	Prospective RCT	On site	7
Esophagus	Squamous cell carcinoma	Detection	NBI > WLI (accuracy: 88.9 vs. 56.5%)	+	Prospective RCT	On site	7
			NBI = iodine CE (accuracy: 95.1 vs. 85.9 %)	+	Prospective non-RCT	On site	24
	Adenocarcinoma	Differential diagnosis	NBI > WLI	+	Prospective RCT	On site	7
		Detection of HGD/early cancer	NBI = WLI + indicocalmine CE	+	Prospective non-RCT	On site	37
		Differential diagnosis	M-NBI > M-WLI	+	Prospective non-RCT	Review	39
		Grade of dysplasia	NBI > WLI	—	Prospective non-RCT	On site	38
		Barrett's esophagus	Detection of SIM	+	Prospective non-RCT	Review	36
Stomach	Adenocarcinoma	Detection	No literature				
		Differential diagnosis of small depressive lesions	C-WLI + M-NBI > M-NBI > C-WLI (accuracy: 96.6 vs. 90.4 vs. 64.8%)	+	Prospective RCT	On site	46
		Differential diagnosis of superficial elevated lesions	M-NBI > C-WLI (accuracy: 92 vs. 74%)	+	Retrospective	Review	57
		Detection of lateral extent of early gastric cancer	M-NBI > indigocalmine CE	+	Prospective RCT	On site	49
		Diagnosis of tumour depth	No pivotal study				
Duodenum	Adenoma	Detection	No pivotal study				
	Adenocarcinoma	Detection	No pivotal study				
		Differential diagnosis	No pivotal study				
	Ampullary tumour	Visualization of tumour margin	NBI > indigocalmine CE	—	Prospective non-RCT	Review	61

NBI: narrow-band imaging; WLI: white light imaging; PPV: positive predictive value; NPV: negative predictive value; SIM: specialized intestinal metaplasia; HGD: high grade dysplasia; Sn.:sensitivity; Sp.: specificity; M-NBI: magnifying NBI; M-WLI: magnifying WLI; CE: chromoendoscopy; C-WLI: conventional nonmagnifying WLI

## Figure Legends

Figure 1. Superficial squamous cell carcinoma in the lower thoracic esophagus.

(A) WLI shows scattered reddish spots in the slightly reddish area.


(B) Lugol chromoendoscopy shows unstained area.

(C) (D) NBI shows clearly defined brownish spots indicating dilated intrapapillary capillary loops.

Figure 2. Microvascular patterns of gastric cancer. (A) Fine network pattern indicates differentiated adenocarcinoma. (B) Corkscrew pattern indicates undifferentiated adenocarcinoma.

Figure 3. White opaque substance (WOS) within an elevated adenoma and well-differentiated adenocarcinoma. (A) The regular distribution of WOS indicates adenoma. (B) The irregular distribution of WOS indicates adenocarcinoma.

Figure 1 (A)



(B)



(C)



(D)







Figure 2 (A)



(B)



Figure 3 (A)



(B)

